

AD-A240 258 NTATION PAGE

Form Approved
OMB No. 0704-0188



* 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, completing the review, and writing the response. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Washington Headquarters Service, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Project Director (0704-0188), Washington, DC 20503.

1. AUTHOR L. R. Goodman		2. REPORT DATE August 1991	3. REPORT TYPE AND DATES COVERED Professional paper
4. TITLE AND SUBTITLE ASSIGNMENT OF SINGLE VALUES TO PROBABILITY INTERVALS, EVALUATION OF CONDITIONAL EVENTS, AND APPLICATIONS TO COMBINATION OF EVIDENCE		5. FUNDING NUMBERS PR: CD32 PE: 0305108K WU: DN488828	
6. AUTHOR(S) L. R. Goodman		8. PERFORMING ORGANIZATION REPORT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Ocean Systems Center San Diego, CA 92152-5000		10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of the Secretary of Defense Research and Engineering Washington, DC 20363		11. SUPPLEMENTARY NOTES	
12a. DISTRIBUTION AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.		12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) A long unrecognized problem in probability and statistics has been the inability to treat inference statements - such as "if b then a" or "a given b" - so that logical combinations of them can be evaluated, compatible with conditional probability. Thus, in the situation where no conditioning occurs - or everything is conditioned on a common antecedent - statements such as "if b then a or if b then c" can be readily addressed with the typical evaluation: $p(a \vee c b) = p_b(a \vee c) = p_b(a) + p_b(c) - p_b(a \wedge c)$, etc., for any probability measure p over the space of events.			
14. SUBJECT TERMS data fusion uncertainty measures combination of evidence game theory		15. NUMBER OF PAGES 16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT SAME AS REPORT

91-10561



Published in *Proceedings of the 28th Annual Bayesian Research Conference*, 1989.

UNCLASSIFIED

1- NAME (Last, first, middle initial) L. R. Goodman	2- TELEPHONE (include Area Code) (619) 553-4014	3- OFFICE SYMBOL Code 421

ASSIGNMENT OF SINGLE VALUES TO PROBABILITY
INTERVALS, EVALUATION OF CONDITIONAL EVENTS,
AND APPLICATIONS TO COMBINATION OF EVIDENCE

DR. I.R. GOODMAN

COMMAND & CONTROL DEPARTMENT,
CODE 421
NAVAL OCEAN SYSTEMS CENTER
SAN DIEGO, CALIFORNIA 92152-5000

*Abstract of paper to be delivered as an invited
lecture to the 28th Annual Bayesian Research
Conference, University of Southern California,
Los Angeles, February 15-16, 1990, under direction of
Prof. Ward Edwards and sponsored by the University
of Southern California.*



Assignment For	
DTIC GMAH	
DTIC TAB	
Unannounced	
Justification	
by	
Distribution	
Availability Codes	
Avail and/or	
Special	
A-1	

ASSIGNMENT OF SINGLE VALUES TO PROBABILITY INTERVALS, EVALUATION OF CONDITIONAL EVENTS AND APPLICATIONS TO COMBINATION OF EVIDENCE

ABSTRACT

A long unrecognized problem in probability and statistics has been the inability to treat inference statements - such as "if b then a" or "a given b" - so that logical combinations of them can be evaluated, compatible with conditional probability. Thus, in the situation where no conditioning occurs - or everything is conditioned on a common antecedent - statements such as "if b then a or if b then c" can be readily addressed with the typical evaluation: $p(a \vee c | b) = p_b(a \vee c) = p_b(a) + p_b(c) - p_b(a \cdot c)$, etc., for any probability measure p over the space of events.

On the other hand, until recently, such simple appearing statements as Δ - "if b then a" and "if a then not d" could not be analyzed within the standard pervue of probability so that one could make sense of the evaluation $p(\Delta)$, compatible with conditional probability, i.e., $p(\text{"if b then a"}) = p(a|b)$ and $p(\text{"if d then not c"}) = p(c'|d)$. (This excludes material implication - and indeed, as Goodman & Nguyen have recently demonstrated [*Conditional Inference and Logic for Intelligent Systems: A Theory of Measure-Free Conditioning*, Chapter 1, North-Holland Press, to appear], no closed operator over a finite boolean algebra of events will also work.) This has lead to the development of a syntactic / algebraic approach to conditioning in probability - much as Boole originally envisioned with his "division" operator, but which was only partially developed by him (although later justified by Hailperin - *Boole's Logic and Probability*) and independently considered from time to time. Only Schay (1958) and, independently, Calabrese (1985), prior to the work here considered, have attempted to develop full-blown conditional event algebras, but their efforts are fraught with empirical and ad hoc components.

In the establishment of such an algebra of conditional events (as in the above reference of Goodman & Nguyen), a program of four parts is required as follows (though not necessary in that order at all times): 1 What algebraic forms, if any, must conditional events take? (Answer: all principal ideal cosets generated from all principal ideal quotient boolean algebras of the original boolean algebra of unconditional events); 2 What functional forms must the conditional event extensions of boolean (unconditional) operators take? (Answer: functional image extensions of all the unconditional point-wise operators of the original boolean algebra to the coset domains); 3 What properties do conditional events and their operators and relations possess? (Answer: Feasible calculus of extended boolean-like operators and partial order extending ordinary subset relations leading to a bounded, distributive, idempotent, DeMorgan, involutive, pseudo-complemented Stone lattice which is also a semi-simple Chang algebra isomorphic to certain variations of Lukasiewicz three-valued logic; and which is a form of Koopman qualitative conditional probability structure, also, which has a full algebraic characterization, extending the Stone Representation Theorem to conditional form); 4 What numerical or semantic properties do these entities possess and what is the nature of assigning a single number - the conditional probability - to a coset of events, which under functional image extensions of probability becomes an interval of numbers in the unit interval? It is this last issue that has not yet been fully satisfactorily addressed.

Given that the assignment is simply $p((a|b)) = p(a|b)$ to the conditional event $(a|b)$, one in effect is attaching a single most representative number in some sense to the interval $\{p(x): x \in (a|b)\} = \{p(x \cdot b') + p(a \cdot b): x \text{ arb. } \in \text{boolean alg.}\} = \text{closed interval } [p(a \cdot b), p(a \cdot b) + 1 - p(b)]$, provided p is non-atomic. With this evaluation, one can show a resulting conditional event probability logic which is sound and complete and monotonically preseving partial order of conditional events, etc. But all of this hinges upon the "natural" interpretation of $p((a|b)) = p(a|b)$. Some characterizations for this relation are presented, including a fixed point weighting representation, a modified Renyi-Aczel property, DeFinetti-Lindley uncertainty game approach, and others. But the basic question remains: Why should s/t be assigned to $[s, 1-t+s]$, or equivalently, $s/(1-t+s)$ to $[s, t]$, for all $0 \leq s \leq t \leq 1$?